

## Improvement of Faba bean (*Vicia fabae* L. var. minor) phosphorus uptake and nitrogen fixation in a Tunisian multi local field test

H. MAAZAoui<sup>1,2,4\*</sup>, J.J. DREVON<sup>3</sup>, B. SIFI<sup>1</sup>.

<sup>1</sup> Agronomic techniques and sciences laboratory, INRAT, Hédi Karray Street, 2080 Ariana, Tunisia.

<sup>2</sup> Faculty of sciences of Bizerte

<sup>3</sup> INRA, UMR Eco&Sols, 2 Place Pierre Viala, 34060 Montpellier, France

<sup>4</sup> Carthage University

\*Corresponding author: maazaouihouda@gmail.com

**Abstract** - The commercial cultivar Bachar of faba bean (*Vicia faba* L. var. minor) was inoculated with a local rhizobial strain, named Mat.11, in twelve farmer fields of Mateur agro-ecosystem in northern Tunisia which is a new agriculture approach in Tunisia. Phosphorus uptake, nitrogen accumulation, field yield and the soil analysis were studied. The results indicate a large spatial variation of the efficiency of the symbiosis rhizobia. Significant differences at  $P < 0.01$  were found among the treatments. Thus, without inoculation treatment suggests the potential of valuable rhizobial strains in some soils in nodulation and plant growth. Nevertheless, the inoculation with Mat.11 increased significantly not only the total plant phosphorus uptake (from 1.46 mg P  $pl^{-1}$  to 4.11 mg P  $pl^{-1}$ ) but also nitrogen fixation (0.6 mg N  $pl^{-1}$  to 3.22% mg N  $pl^{-1}$ ) comparing to without inoculation treatment in many sites. The inoculation has also increased the grain yield comparing to the control treatment. It is concluded that faba bean inoculation with Mat11 improve significantly nodulation and plant growth. Further investigations should address the variation SNF among local genotypes and the symbiosis mechanisms to stabilize the legume yield in soils with low fertilizer input.

**Keywords:** Faba bean, nitrogen, phosphorus, rhizobia, symbiosis.

### 1. Introduction

Nitrogen nutrition legume is even more difficult to study due to the combination of two very different ways, the nitrogen fixation Atmospheric and assimilation of mineral nitrogen. The contribution of each of these two paths to the nitrogen supply of the plant varies considerably depending on the species and the conditions of culture (Wery 1987). The improvement of the production of pulses and their use as previous Cultural therefore requires better understand the relationships between the assimilation and nitrogen fixation on the one hand, and the influence of environmental factors and techniques cultivation on these two channels, on the other hand.

In northern Tunisia, forests and range lands in areas with acid soils and annual rainfall as high as 1200 mm are actually converted into annual cropping of cereals and legumes. The symbiosis between legumes and rhizobia is the main source of biologically fixed nitrogen for agricultural systems. Estimates for field grown legumes reveal that up to 80 per cent of the legume N demand is met by symbiotic  $N_2$  fixation (SNF) (Graham & Vance 2003; Vance 2001). Total global  $N_2$  fixation from SNF has been estimated to 100–290 million tons N  $year^{-1}$ , compared with 83 million tons N fixed industrially in fertilizer production (Herridge 2008). Moreover, SNF is potentially beneficial for subsequent crops in rotations (Gan et al. 2003) and to non-legumes in mixed stands. Also, inclusion of legumes in crop rotation was found to improve soil physical and chemical properties (Biederbeck et al. 2005). Legumes in cropping systems were also found to be useful in controlling pests, diseases, and weeds (Howieson et al. 2000). Among grain legumes, faba bean (*Vicia faba*) is the fourth most important pulse crop in the world. It occupies the lowest area planted to legume crops and cereal monoculture in Arab countries especially in Tunisia. Faba bean is commonly grown in northern Tunisia and covers a surface area of about 56,600 ha (Ministry of agriculture 2011). In addition, the soil reaction affects the survival and persistence of rhizobia and their nodulation on roots, and therefore also the ability of faba bean plants to fix aerial nitrogen.

Phosphorus is an essential element for the growth and development of plants Phosphorus deficiency has previously been reported to decrease nodule biomass more than host growth in soybean (Drevon & Hatwig 1997). However, in agricultural soils, the major portion of phosphorus is insoluble and therefore inaccessible to plants. In the calcareous soil of the Mediterranean basin, symbiotic nitrogen fixation (SNF) is frequently limited due to low P nutrient availability (Broughton et al. 2003). Phosphorous deficiency affects legumes more than other crops since SNF is an energetically expensive process requiring more inorganic P (Pi) than mineral nitrogen N (Vadez 2001). The P deficiency affects symbiotic fixation of N<sub>2</sub> by limiting growth and survival of rhizobia (O' Hara et al. 1988), nodule formation (Latati et al. 2014) and functioning (Tang et al. 2001a), and host plant growth (Tsvetkova and Gorgiev 2003; Latati et al. 2015). To cope with this low P availability we suggest that rhizobia inoculation could be an alternative practice that is widely adopted in tunisian agriculture so far. Thus, this multilocal field study aimed to highlight whether the rhizobia inoculation might alleviate the negative effect of P disponibility and improves plant growth and nutrition without additional nitrogen in northern Tunisian soils. The current study was designed to evaluate the response of faba bean to local rhizobial inoculation in terms of growth parameters, nitrogen fixation and yield in Mateur multi-location field tests.

## 2. Materials and methods

### 2.1. Multi located field trials

The study was conducted during 2012-2013 in 12 farmer's field sites with mean distances of 20 km between each site at Mateur, eastern north of Tunisia (Latitude: 37°02'N, Longitude: 9°39'E and altitude 36 m). The mean annual temperature across the sites ranged between 10 and 26°C, and the mean annual rain fall ranged between 450 and 600 mm with maximum rainfall in the period between November and February. The lower valley of Medjerda's Mediterranean climate is characterized by cold winters and dry summer. Soils of the experimental sites were characterized by a standard sampling of the top 30 cm using an auger at sowing stage. Table 1 shows physical and chemical soil properties for the experimental sites.

**Table 1** Mat strain inoculation effect on nitrogen and phosphorus content of faba bean (*Vicia fabae*.M)

Sites	Nitrogen content mg/pl		Phosphorus content mg/pl	
	Without inoculation	With inoculation	Without inoculation	With inoculation
A9	0,54a	<b>1,07de</b>	1,95ef	2,35e
A5	2,04c	1,63d	2,92f	<b>2,68cd</b>
A8	1,34d	<b>2,57c</b>	2,11f	<b>2,87c</b>
A12	0,75e	1,38d	1,72g	<b>2,57e</b>
A4	6,19a	2,62c	2,33ef	2,65e
A6	1,11de	1,58d	1,52g	2,2ef
A2	1,11e	1e	1,56g	<b>3,22b</b>
A10	2,9bc	3,22b	1,66g	1,46g
A7	0,95de	<b>1,79d</b>	1,91g	<b>3,42b</b>
A11	0,68e	0,69e	2,46c	3,21b
A3	0,47e	<b>0,7e</b>	3,61ab	2,62c
A1	0,89e	0,79e	2,54c	<b>4,11a</b>
F value	37.81**	6 8.14**	27.81**	67.05**

### 2.2. Biological material and treatments

Two treatments, each on elementary plot of 100 m<sup>2</sup>, were compared in twelve field site: non inoculated (control) and inoculated with Mat strain (Mat.11) containing ten rows distant of 50 cm at a sowing density of 30 seeds per m and cultivated with Bachar variety selected by INRAT. In order to prepare fresh inoculum containing the same number of bacterial population for all rhizobia under study, a colony of Mateur strain was transferred into Erlenmeyer containing 100 mL of Yeast extract mannitol medium (Vincent 1970). Inoculated flasks were incubated at 27°C on a rotary shaker (150 rpm) for 72 hours. All bacterial suspensions were adjusted to about 10<sup>8</sup> CFU ml<sup>-1</sup>. This suspension was used to inoculate the emerging seedlings using a watering can. Plants were irrigated by rainfall regime.

**Table 2** Mat Strain inoculation effect on 100g seed dry weight and yield

	yield kg/Ha		100g seed dry weight	
	Without inoculation	With inoculation	Without inoculation	With inoculation
A9	4031,6	2296,4	36b	42a
A5	2003,2	1737,5	44,2a	32b
A8	2839,1	4203,3	40b	47a
A12	2154,4	2476,1	29b	32b
A4	1464,1	2267,5	48a	38b
A6	3327,6	3294,3	37a	34a
A2	2655,9	4347	45b	49b
A10	2555,5	2748,9	28a	30a
A7	4068	4306,1	44a	45a
A11	4386,6	3298,1	37a	34a
A3	1543,2	1772,6	34a	39a
A1	1721,2	2505,3	31a	38a
F value			9.43**	8.37**

Mean in each column followed by the same letters are not significantly different at  $P < 0.05$  according to Duncan's multiple range test. \*: Significant at  $P < 0.05$ ; \*\*: Significant at  $P < 0.01$

### 2.3. Plant parameters

Ten plants were harvested at early flowering stage of each treatment at a depth of 20 cm and separated into shoot root and nodules.

For N determination, the sample was digested in hot concentrated  $H_2SO_4$ , as described by Kjeldahl. Soil P availability was determined after extraction in  $NaHCO_3$  (Olsen et al. 1954).

For P determination, the plants were performed according to standardized by AFNOR method (1969). The method is to pass in solution, the plant sample previously introduced in mini reactors and mineralized microwave in under the attack of concentrated nitric acid to pressures of the order of 110 bars and a temperature up to  $180^\circ C$  for 15 minutes. Quantification of anhydride phosphoric acid solution was carried pair reagent colorimetric assay vanadomolybdic revealing yellow units whose OD was read to 460 nm.


### 2.4. Statistical analysis

All values were means of 20 replicates. Data were analysed statistically by running ANOVA of Statistical Package for the Social Sciences (SPSS) version 20.0 (SPSS Inc., Chicago, USA). The mean values were compared using Duncan's multiple range test ( $P < 0.05$ ).

## 3. Results and discussion

### 3.1. Nitrogen and phosphorus uptake.


In order to assess the nitrogen fixation per site, the values of total nitrogen plant were plotted in figure 1. Total nitrogen varied from  $1.0 \text{ mg N pl}^{-1}$  in A4 to  $0.5 \text{ mg N pl}^{-1}$  in A2 without inoculation, and from  $1.7 \text{ mg N pl}^{-1}$  in A8 to  $0.5 \text{ mg N pl}^{-1}$  in A4 with inoculation. Overall, the inoculation with local Mat strain increased the mean total N content in four sites A9 ( $1.07^* \text{ mg N pl}^{-1}$ ), A8 ( $2.05^{**} \text{ mg N pl}^{-1}$ ), A7 ( $1.79 \text{ mg N pl}^{-1}$ ) and A3 ( $0.7^* \text{ mg N pl}^{-1}$ )



**Figure.1** Effect of rhizobial inoculation on nitrogen content of cultivated faba bean in field sites. Data are means and SD of 20 replicates harvested at flowering stage without (□) or with inoculation (■).

Faba bean inoculation increased plant P uptake from 5per cent in A9 to 30per cent in A12, 150per cent in A1 and A7 sites. The mean P plant value varied, from 1.46 mg P pl<sup>-1</sup> in A10 to (4.11\*\*) mg P pl<sup>-1</sup> in A1 farmer field A2 (3.22\*\* mg P pl<sup>-1</sup>), A8 (2.85\*mg P pl<sup>-1</sup>), A7 (3.42\*\* mg P pl<sup>-1</sup>) and A12 (2.47\* mg P pl<sup>-1</sup>). Whereas, in control plots, plant phosphorus uptake varied from 1.36 mg P pl<sup>-1</sup> in A2 to 3.61 mg P pl<sup>-1</sup> in A3 (fig.2). The inoculation with rhizobia strain had no effect on plant P uptake in A3. The highest Plant P uptake was observed in A1 with inoculation, and the lowest P plant uptake root biomass was observed in A2 and A6 without inoculation. The plant P uptake is not related with soil P content (Table 3 and fig2). Phosphorus is generally deficient and limits the potential input of SNF (Waluyo et al. 2004) by reducing the number and biomass of nodules and their nitrogenase activity (Ribet & Drevon 1995); (Vadez et al. 1996; Qiao et al. 2007). However in our observations, P did not affect the nodulation whatever the site. Nevertheless, rhizobial inoculation increased plant P uptake in almost 40per cent of the experimental sites. This finding is in agreement with results showing that increasing the P levels in the plant should also increase their allocation to the various organs and thus improve the quantity and quality of seeds (Lagrange 2009).

The high nitrogen levels were recorded in plants inoculated and fertilized with nitrogen while the lowest were recorded in control plants. The results allowed inferring that the inoculation and fertilization have a significant effect on the nitrogen accumulation. The phosphorus intake among inoculated plants has a significant effect on the nitrogen content. Compared with other Fabaceae, (Zamen-Alah 2007) confirmed the high sensitivity of the symbiotic nitrogen fixation in phosphate fertilization in bean. Moreover, some studies have shown that the metabolism of symbiotic nitrogen fixation requires more phosphorus than the growth of the aerial part (Gundwana et al. 1992). According to Bargaz et al. (2011), the efficiency of phosphorus use in nitrogen-fixing symbioses is closely linked to adequate distribution of P between the aerial parts and roots nodulated and between the nodules and roots.




**Figure.2** Effect of rhizobial inoculation on plant phosphorus of cultivated faba bean in field sites. Data are means and SD of 20 replicates harvested at flowering stage without (□) or with inoculation (■).

**Table 3:** Chemical characteristic of the differents fields location at depth of 0-20cm

samples	Sand for100 g soil	Clay	Silt	Total Calcareous	Activ Calcareous	Organic Matter	pH	CE mmhos/cm	N %	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
A9	10	60	30	3	2	3,59	7,61	0,15	1,52	41	260
A5	20	25	55	32	11	3,48	7,63	0,14	1,80	94	380
A8	25	35	40	47	18	2,32	7,26	0,13	1,80	83	170
A12	30	30	40	31	5	2,64	7,15	0,17	1,84	67	270
A4	10	40	50	30	9	2,64	7,55	0,15	1,60	55	230
A6	15	35	50	31	13	1,79	7,67	0,15	1,54	38	240
A2	40	35	25	32	9	3,69	7,30	0,17	1,96	89	310
A10	10	25	65	27	8	1,79	7,76	0,14	1,80	63	320
A7	15	35	50	15	6	1,79	7,09	0,14	1,80	52	400
A11	25	30	45	24	9	2,64	7,08	0,14	1,80	97	130
A3	20	45	35	15	6	2,32	7,60	0,14	1,64	67	310
A1	15	35	50	18	6	2,85	7,20	0,14	1,84	80	370


### 3.2. Effect of inoculation on faba bean yield

In all control field mean yield varied among the site between 1300 kg/ha in A4 and 4300 kg/ha in A11 (Fig.3). The mean yield of faba bean inoculation with Mat.11 varied from 1600 kg/ha in A3 to 4300 kg/ha in A2, A8 and A7. Inoculation increased yield in all sites trials except A9, A5 and A11 where the effect was negative. Several studies have reported a positive effect of inoculation leading to a significant improvement in seed yield (Karasu et al. 2009). Although it is a small contribution to crop production.



**Figure.3** Effect of rhizobial inoculation on yield of cultivated faba bean in field sites. Data are means and SD of 20 replicates harvested at flowering stage without (□) or with inoculation (■)

Quality of faba bean grain was improved by inoculation with Mat strain. The mean of 100 grain dry weight in control plants varied from 28 g in A3 to 1.0 g in A10 (Fig.4). With inoculation of faba bean with Mat.11 32 g in A12 to 49 g in A2. Inoculation decreased 100 grain dry weight in all sites trials except A5 and A9. Generally, a small increase in 100 grain dry weight of the order of 5 to 15 % is observed after inoculation with Mat strain (Sindhu et al. 2010). However, the desired impact of bio fertilizer on legume is obtained under some field conditions.



**Figure.4** Effect of rhizobial inoculation on 100 seed dry weight (D) of cultivated faba bean in field sites. Data are means and SD of 20 replicates harvested at flowering stage without (□) or with inoculation (■).

### 4. Conclusion

Strain used to inoculate a plot increased the total phosphorus and total nitrogen content of plants for the majority of the sites which results in improved yields. The amount of nitrogen fixed by symbiotic way remains very useful in maintaining and restoring soil fertility and in increased yield compared to non-inoculated cultures. It is also important to note that, in addition to the significant effect of inoculation on the yield of legumes, nitrogen application at planting is precious in agro systems. Thus further comparisons of faba bean sites among the diversity of the producers and fields of the region may be fruitful to assess the transferability of the rhizobial-inoculation. Further exploration of the interaction between the native rhizobia and locally-used cultivars faba bean may be recommended for optimization of this inoculant-biotechnology.

**Acknowledgments** This work was supported by the Great Federative Project of Agropolis named FABATROPIMED under the reference ID 1001-009 and framework of Tunisia-French cooperation AUF-PCSI 63113PS012 for the scholar internship of Maazaoui Houda in Montpellier.



## 5. References

- Bargaz A, Drevon JJ, Oufdou K, Mandri B, Faghire M, Ghoulam CH (2011)** Nodule phosphorus requirement and O<sub>2</sub> uptake in common bean genotypes under phosphorus deficiency. *Soil and Plant Science*. 2011:1-10.
- Biederbeck VO, Zentner RP, Campbell CA (2005)** Soil microbial populations and activities as influenced by legume green fallow in a semiarid climate. *Soil Biology and Biochemistry* 37, 1775-1784.
- Broughton JW, Hernandez G, Blair M, Beebe S, Gepts P, Van derleyden J (2003)**. Beans (*Phaseolus* spp) model food legumes. *Plant and Soil* 252, 55-128.
- DGPA (2011)** Followed report and evaluation of field crops. Ministry of agriculture (2011)
- Drevon JJ, Hartwig UA (1997)** Phosphorus deficiency increases the argon-induced decline of nodule nitrogenase activity in soy bean and alfalfa, *Planta* 201, 463-469.
- Gunawardena SF, Danso SK, Zapata F (1992)** Phosphorus requirement and sources of nitrogen in three soybean (*Glycine max*) genotypes: Bragg, nts 382 and Chippewa. *Plant and soil*, 151:19-26
- Gan YT, Miller PR, Mc Conkey BG, Zentner, RP, Stevenon FC, Mc Donald, CL (2003)** Influence of diverse cropping sequences on durum wheat yield and protein in the semiarid northern Great Plains. *Agronomy Journal* 95, 245-252.
- Herridge DF, Peoples MB, Boddey RM (2008)** Global inputs of biological nitrogen fixation in agricultural systems. *Plant and Soil* 311, 1-18.
- Howieson JG, O'Hara GW, Carr SJ (2000)** Changing roles for legumes in Mediterranean agriculture: developments from an Australian perspective. *Field Crops Research* 65, 107-122.
- Latati Mourad, Blavet Didier, Alkama Nora, Laoufi H, Drevon Jean-Jacques, Gérard Férédéric, Pansu Marc, Ounane Sidi Mohamed (2014)** The intercropping cowpea-maize improves soil phosphorus availability and maize yields in an alkaline soil. *Plant Soil* 385:181–191.
- Latati Mourad, Adnane Bargaz, Baroudi Belarbia, Mohamed Lazali, Samia Benlahrech, Siham Tellah, Ghiles Kaci, Jean Jacques Drevon, Sidi Mohamed Ounane (2015)** The intercropping common bean with maize improves the rhizobial efficiency, resource use and grain yield under low phosphorus availability. *European Journal Of Agronomy*, Volume 72, January 2016, page 80-90.
- Lagrange (2009)** Ecological and microbiological studies of species of *Costularia* (Cyperaceae), pioneers of ultramafic soils in New Caledonia: application to ecological restoration.
- O'Hara GW, Bunkered N, Dilworth MJ (1988)** Mineral constraints to nitrogen fixation, *Plant Soil* 108, 93-110.
- Olsen SR, Cole CV, Watanabe FS, Dean LA (1954)** Estimation of Available Phosphorus in Soil by Extraction with Sodium Bicarbonate; Circular 939. USDA, Washington DC, USA, pp. p19.
- Qiao YF, Tang C, Han XZ, Miao SJ (2007)** Phosphorus deficiency delays the onset of nodule function in soybean (*Glycine max* Murr.). *Journal Plant Nutrition* 30:1341–1353.
- Ribet J, Drevon JJ (1995)** Increase in permeability to oxygen and in oxygen uptake of soybean nodules under limiting phosphorus nutrition. *Physiol Plant* 94:298–304.
- Sindhu SS, Dua S, Verma MK, Khandewal A (2010)** Growth Promotion of Legumes by Inoculation of Rhizosphere Bacteria. M.S.Khan et al. (eds.). *Microbes for Legume Improvement*. Springer-Verlag Wien, Germany. DOI 10.1007/978-3-211-99753-6..
- Tang C, Hinsinger P, Drevon JJ, Jaillard B (2001a)** Phosphorus deficiency impairs early nodule functioning and enhances proton release in roots of *Medicago truncatula* L., *Ann. Bot* 88, 131-138.
- Tsvetkova GE, Georgiev GI (2003)** Effect of phosphorus nutrition on the nodulation, nitrogen fixation and nutrient –use efficiency of *Bradyrhizobium japonicum* soybean (*Glycine max* L. Merr.) symbiosis, *Bulg.J. Plant Physiol. Special issue* 331-335.
- Vadez V, Rodier F, Payre H, Drevon JJ (1996)** Nodule permeability to O<sub>2</sub> and nitrogenase linked respiration in bean genotypes varying in the tolerance of N<sub>2</sub> fixation to P deficiency. *Plant Physiol Biochemistry* 34:871–878.
- Vadez, Drevon JJ (2001)** Genotypic variability in p use efficiency for symbiotic n<sub>2</sub> fixation in common bean (*phaseolus vulgaris* l). *Agronomie* 21 (2001) 691-699.
- Vance CP (2001)** Symbiotic nitrogen fixation and phosphorus acquisition. *Plant nutrition in a world of declining renewable resources*. *Plant Physiology* 127, 390-397.
- Waluyo SH, Lie TA, Mannetje L (2004)** Effect of phosphate on nodule primordia of soybean (*Glycine max*) in acid soils in rhizotron experiments. *Indonesian J Agric Scil* 5:27–44.
- Wery J (1987)**. Relation entre la nutrition azotée et la production chez les légumineuses. In : Actes du colloque, Nutrition azotée des légumineuses, 19-21 novembre 1985, Versailles, France. Paris : INRA.
- Zamen Allah M, (2007)**. Effects of inoculation with rhizobia strains on tolerance of different bean cultivars (*Phaseolus vulgaris* L.) to salt stress and phosphorus limitation. Thesis, p: 9.